

TECHNICAL BULLETIN

PURELL[®] Advanced Instant Hand Sanitizer Technical Data

INDICATIONS: Hand sanitizer to help reduce bacteria on the skin that could cause disease. Recommended for repeated use.

DIRECTIONS: Place enough product in your palm to thoroughly cover your hands. Rub hands together briskly until dry. Children under 6 years of age should be supervised when using this product.

Physical Properties

Appearance: **Clear liquid**

Fragrance: **Citrus fragrance**

Form: **Gel**

pH: **6.5 – 8.5**

INCI Name*
Active:
Ethyl alcohol 70% v/v
Also Contains:
Water (Aqua)
Isopropyl Alcohol
Caprylyl Glycol
Glycerin
Isopropyl Myristate
Tocopheryl Acetate
Acrylates/C10-30 Alkyl Acrylate Crosspolymer
Aminomethyl Propanol
Fragrance (Parfum)

*International Nomenclature Cosmetic Ingredient

Efficacy Data – *In Vivo*

Healthcare Personnel Handwash

Objective: This study evaluated the antimicrobial effectiveness of one (1) test product and one (1) control product using a Health-Care Personnel Handwash Procedure, as per methodology specified by the Food and Drug Administration (FR 59:116, 17 Jun 94).

Description of Test: Twenty-four (24) subjects utilized test product and twenty-seven (27) utilized the positive control reference product (51 total). The antimicrobial effectiveness of test product and control product for use as Health-Care Personnel Handwashes were determined using eleven (11) consecutive hand contaminations, the first followed by a sample for baseline, and the remaining ten (10) by product applications. Microbial samples were taken at baseline and after product applications one (1), three (3), seven (7), and ten (10). All sampling of the hands was performed using the Glove Juice Sampling Procedure. *Serratia marcescens* (ATCC #14756) was the marker organism used for hand contaminations. The FDA requires products to achieve a minimum 2 log₁₀ reduction after one application and 3 log₁₀ reduction after 10 applications.

Independent Laboratory; Study# BioScience Laboratories, Inc., Bozeman, MT; 100635-101

Date: 29 October 2010

Results:

Application Number	Test Product Log ₁₀ Reduction	Control Product Log ₁₀ Reduction
1	3.20	3.05
10	3.60	4.76

Conclusions: Test product meets FDA Healthcare Personnel Handwash requirements when 2 mL of product is applied to the hands and rubbed in until dry.

Healthcare Personnel Handwash

Objective: This study evaluated the antimicrobial effectiveness of one (1) test product and one (1) control product using a Health-Care Personnel Handwash Procedure, as per methodology specified by the Food and Drug Administration (FR 59:116, 17 Jun 94).

Description of Test: Twenty-six (26) subjects utilized test product and twenty-four (24) utilized the positive control reference product (50 total). The antimicrobial effectiveness of test product and control product for use as Health-Care Personnel Handwashes were determined using eleven (11) consecutive hand contaminations, the first followed by a sample for baseline, and the remaining ten (10) by product applications. Microbial samples were taken at baseline and after product applications one (1), three (3), seven (7), and ten (10). All sampling of the hands was performed using the Glove Juice Sampling Procedure. *Serratia marcescens* (ATCC #14756) was the marker organism used for hand contaminations. The FDA requires products to achieve a minimum 2 log₁₀ reduction after one application and 3 log₁₀ reduction after 10 applications.

Independent Laboratory; Study# BioScience Laboratories, Inc., Bozeman, MT; 111016-101

Date: 19 March 2012

Results:

Application Number	Test Product Log ₁₀ Reduction	Control Product Log ₁₀ Reduction
1	2.85	2.76
10	3.28	4.50

Conclusions: Test product meets FDA Healthcare Personnel Handwash requirements when 1.1 mL of product is applied to the hands and rubbed in until dry.

Efficacy Data – *In Vitro*

Timed – Exposure Kill Evaluation

Objective: Evaluate the antimicrobial effectiveness of the product *in vitro*.

Description of Test: Fifteen (15) second exposure kill evaluations were performed utilizing fifty-six (56) challenge microorganism strains. The challenge inoculum was introduced to the test product at time zero; a portion of the sample was removed and placed in neutralizing media at the appropriate time (15 seconds). Standard plate counting techniques were

used to enumerate viable challenge microorganisms.

Independent Laboratory; **BioScience Laboratories, Inc., Bozeman, MT, USA; 091106-201**

Date: **19 October 2010**

Results:

Challenge Microbe	ATCC No.	Exposure (seconds)	Percent Reduction
<i>Acinetobacter baumannii</i>	19606	15	99.9999
<i>Bacteroides fragilis</i>	25285	15	99.9913
<i>Burkholderia cepacia</i>	25416	15	99.9999
<i>Burkholderia cepacia</i>	25608	15	99.9999
<i>Campylobacter jejuni</i>	29428	15	99.9999
<i>Citrobacter freundii</i>	8090	15	99.9999
<i>Clostridium difficile</i> (vegetative cells)	9689	15	99.9943
<i>Clostridium perfringens</i> (vegetative cells)	13124	15	99.9999
<i>Corynebacterium diphtheriae</i>	11913	15	99.9999
<i>Enterobacter aerogenes</i>	13048	15	99.9999
<i>Enterococcus faecalis</i>	19433	15	99.9999
<i>Enterococcus faecalis</i>	29212	15	99.9999
<i>Enterococcus faecalis</i> VRE	51299	15	99.9999
<i>Enterococcus faecalis</i> VRE	51575	15	99.9999
<i>Enterococcus faecium</i>	19434	15	99.9999
<i>Enterococcus faecium</i> (MDR, VRE)	51559	15	99.9999
<i>Escherichia coli</i>	11775	15	99.9999
<i>Escherichia coli</i>	25922	15	99.9999
<i>Escherichia coli</i> (O157:H7)	43888	15	99.9999
<i>Escherichia coli</i> (MDR, ESBL)	BAA-196	15	99.9999
<i>Escherichia coli</i> ESBL; Carbapenemase-Producing	BSLI #082710EcC P1*	15	99.9998
<i>Haemophilus influenzae</i> MDR	33930	15	99.9999
<i>Klebsiella pneumonia</i> Ozaenae	11296	15	99.9999
<i>Klebsiella pneumonia pneumoniae</i>	13883	15	99.9998
<i>Klebsiella pneumoniae pneumoniae</i>	27736	15	99.9998
<i>Klebsiella pneumonia</i> KPC 2 Positive; Carbapenemase Producing	BSLI#081710 KPCI*	15	99.9998
<i>Lactobacillus plantarum</i>	14917	15	99.9999
<i>Listeria monocytogenes</i>	7644	15	99.9999
<i>Micrococcus luteus</i>	7468	15	99.9992
<i>Proteus hauseri</i>	13315	15	99.9999
<i>Proteus mirabilis</i>	7002	15	99.9999
<i>Pseudomonas aeruginosa</i>	15442	15	99.9999
<i>Pseudomonas aeruginosa</i>	27853	15	99.9999
<i>Salmonella enterica enterica</i> serovar <i>Enteritidis</i>	13076	15	99.9999
<i>Serratia marcescens</i>	8100	15	99.9999
<i>Serratia marcescens</i>	14756	15	99.9999
<i>Shigella dysenteriae</i>	13313	15	99.9999
<i>Shigella sonnei</i>	11060	15	99.9999
<i>Staphylococcus aureus aureus</i>	6538	15	99.9999

<i>Staphylococcus aureus aureus</i>	29213	15	99.9999
<i>Staphylococcus aureus aureus (MRSA)</i>	33591	15	99.9999
<i>Staphylococcus aureus aureus (MRSA)</i>	33592	15	99.9999
<i>Staphylococcus aureus (MRSA) (VRSA)</i>	062707 NARSAVRSal	15	99.9999
<i>Staphylococcus aureus (MRSA) (NARSA Strain NRS384;USA 300)</i>	12085 NRSa384	15	99.9999
<i>Staphylococcus epidermidis</i>	12228	15	99.9998
<i>Staphylococcus epidermidis MRSE</i>	51625	15	99.9998
<i>Staphylococcus haemolyticus</i>	43252	15	99.9998
<i>Staphylococcus hominis hominis</i>	27845	15	99.9997
<i>Staphylococcus saprophyticus</i>	49453	15	99.9999
<i>Streptococcus pneumoniae</i>	6303	15	99.9999
<i>Streptococcus pneumoniae</i>	49619	15	99.9999
<i>Streptococcus pyogenes</i>	14289	15	99.9999
<i>Streptococcus pyogenes</i>	19615	15	99.9999
Yeasts	ATCC No.	Exposure (seconds)	Percent Reduction
<i>Candida albicans</i>	18804	15	99.9999
<i>Candida albicans</i>	66027	15	99.9999
<i>Candida tropicalis</i>	13803	15	99.9999

Conclusions:

Very effective reduction of gram-negative and gram-positive bacteria and yeasts was demonstrated.

ESBL- Extended Spectrum Beta-Lactamase Producer

MDR – Multi-Drug Resistant

MRSA - Methicillin Resistant *Staphylococcus aureus*

MRSE – Methicillin Resistant *Staphylococcus epidermidis*

NARSA – Network on the Antimicrobial Resistance in *Staphylococcus aureus*

VRE – Vancomycin-Resistant *Enterococcus*

* - Clinical Isolate

Irritancy Data and Allergy Test Results

21 Day Cumulative Irritancy Assay with Delayed Challenge

Objective:	Evaluation of skin irritation potential in humans.
Description of Test:	Phillips et al (Toxic and Applied Pharmacology 21:369-382) summarizes the method utilized for this evaluation. Fresh materials are applied daily, 5 days per week, for 21 days to the same site (patches were not moved or reapplied on the weekends).
Independent Laboratory; Study#	RCTS, INC. Irving, TX, USA; 2749
Date:	6 October 2010
Results:	CIT Average Score = 0.35 (scale 0 – 4; Baby Oil = 0.24) Challenge Phase: Non-sensitizing
Conclusions:	Product has a low potential for skin irritation and allergic contact dermatitis.

Human Repeated Insult Patch Test

Objective:	To determine the irritation and sensitization (contact allergy) potential of a test material after repeated application to the skin of subjects.
Description of Test:	<p>This study was conducted utilizing a standard protocol and a total of fifty-two (52) subjects.</p> <p>Subjects were requested to bathe or wash as usual before arrival at the facility. Patches containing the test material were then affixed directly to the skin of the intrascapular regions of the back, to the right or left of the midline and subjects were dismissed with instructions not to wet or expose the test area to direct sunlight.</p> <p>Subjects were instructed to remove the patches approximately 48 hours after the first application and 24 hours thereafter for the remainder of the study. This procedure was repeated until a series of nine (9) consecutive, 24-hour exposures had been made three (3) times a week for three (3) consecutive weeks. Prior to each reapplication, the test sites were evaluated by trained laboratory personnel.</p> <p>Following a 10-14 day rest period a retest/challenge dose was applied once to a previously unexposed test site. Test sites were evaluated by trained laboratory personnel 48 and 96 hours after application.</p> <p>In the event of an adverse reaction, the area of erythema and edema were measured. Edema is estimated by the</p>

evaluation of the skin with respect to the contour of the unaffected normal skin.
Subjects were instructed to report any delayed reactions that might occur after the final reading.

Independent Laboratory; Study#: BioScreen Testing Services
Torrance, CA, USA; 10-114A

Date: 17 September 2010

Results: No observed dermal reactions.

Conclusions: No demonstrated potential for eliciting dermal irritation or sensitization.

Compatibility Test Results

Compatibility Study To Measure The Effects Of PURELL Advanced Instant Hand Sanitizer On The Antimicrobial Properties Of A Chlorhexidine Gluconate Surgical Scrub Formulation

Objective: Assess the compatibility of the test article with a known Chlorhexidine Gluconate (CHG) Surgical Scrub using a pig skin procedure.

Description of Test: *Serratia marcescens* ATCC 14756 was used as the indicator organism. The inoculum was applied to sterilized, prepared pigskins and allowed to dry. For baseline samples, skins incubated at room temperature for 2 hours prior to sampling. For the positive control (4% CHG product alone), and test samples (test product applied either before or after a 4% CHG product), products were applied to dried skins then allowed to incubate for 2 hours prior to sampling. Dilutions and plating was done utilizing standard microbiological techniques. Log₁₀ reductions from baseline were calculated and statistical analysis was conducted to determine whether statistical differences exist between the positive control and the test product samples. A product is considered CHG compatible if the log reduction for the test product in combination with 4% CHG product is not significantly inferior to the positive control.

Independent Laboratory; Study#: BioScience Laboratories, Bozeman, MT, USA; 110112-250

Date: 19 April 2011

Conclusion The log reduction of the test product used before or after the CHG product is not significantly different than the log reduction of the CHG product when used alone. Therefore, PURELL Advanced Instant Hand Sanitizer does not interfere with the antimicrobial efficacy of CHG and is compatible with CHG containing products.

Glove Compatibility

Test Method	ASTM D5151-99 Glove samples were immersed in product for a period of 2 hours and then examined for leaks. The control samples were not exposed to product.
Testing Lab; Study#:	Smithers RAPRA Inc., Akron, OH, USA; F19864JE
Date	25 April 2011
Purpose of Study	Determine the effect of product on Medical Gloves including latex, nitrile and vinyl gloves.
Sample Size:	100 control gloves and 100 gloves were tested with PURELL® Advanced Instant Hand Sanitizer on each of three glove types. Tested were Latex, Vinyl and Nitrile gloves.
Results:	Latex, nitrile, and vinyl gloves exposed to product were not significantly different than the control gloves.
Summary:	The test product did not significantly impact the integrity of latex, nitrile and vinyl medical gloves.